



Flea Treatment Preparation

Before Service Can Be Performed

Service Date: _____ **Service Time:** _____

1. Remove all pets, including birds and fish, from your home. If the fish tanks cannot be easily removed from the house, unplug the air pump and cover the tank with a damp cloth. In those instances where turning off the pump may be injurious to fish (e.g., salt water fish), it will be the customer's responsibility to remove the fish. Pestgon is not responsible for any animals, including birds and fish, left in the home. Any exotic pets or animals may need to be kept out for longer periods of time (e.g., tarantulas, reptiles, etc.)
2. Do not water for 24 hours. Pick up all toys inside and outside the house.
3. Animals must be treated for fleas the same day service is to be performed.
4. Vacuum all carpeted areas, all upholstered furniture (be sure to vacuum under and in between the cushions) and any furniture upon which pets sleep or rest. Animal bedding should be washed in hot water or discarded. The vacuum bag must be discarded outside the house.
5. Where possible, remove items from the floor (bean bag chairs, small plants, boxes, etc.) Also remove items from underneath beds and furniture (especially toys and clothing). Remove pet dishes from the floor.
6. Remove items from floors of closets to allow the carpeted surface to be treated. Do not store these items on fabric furniture, as these will also need to be treated.
7. Remove all open food products from counter tops and place in sealed containers or in the refrigerator.
8. All people and pets must be out of the house 4 hours, or until the spray is dry. The yard should be vacated until dry. Birds and small or exotic pets should be kept away for 24 hours. Persons allergic to pesticides, those with respiratory problems and those who are pregnant should check with a physician about the proper re-entry time.
9. Upon returning, windows may be opened.
10. Wait three weeks to allow the chemical to work before expecting satisfactory results.
11. Vacuum area thoroughly 24 to 48 hours after treatment and then every three days for the next three weeks