

NET CONTENTS: 4 LBS.

OF CHILDREN DANGER

See back panel for additional Precautionary Statements



FIRST AID

IF SWALLOWED: CALL PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY. If less than ten (10) minutes have passed since poison was taken, give 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Have patient lie down in quiet, darkened room and keep warm and quiet. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air, If not breathing, give artificial respiration, Get medical attention.

IF IN EYES: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water, Get medical attention.

NOTE TO PHYSICIAN: (A) Administer 100% oxygen by positive pressure to provide as much pulmonary gas exchange as possible despite seizures, (B) Administer ANTICONVULSANT DRUGS intravenously to control convulsions. NOTICE: It may be difficult or impossible to stop the seizure activity without stopping respiration. Be prepared to maintain pulmonary ventilation mechanically. Tracheotomy may be necessary if seizures are prolonged.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of a medical emergency call SafetyCall* at 1-866-897-8050.